

## A-level Chemistry (7405/3)

### Paper 3

Specimen 2014

Session

2 hours

#### Materials

For this paper you must have:

- the Data Booklet, provided as an insert
- a ruler
- a calculator.

#### Instructions

- Answer **all** questions.
- Show **all** your working.

#### Information

- The maximum mark for this paper is 90.

Please write clearly, in block capitals, to allow character computer recognition.

Centre number

Candidate number

Surname

Forename(s)

Candidate signature \_\_\_\_\_

**Section A**

Answer **all** questions in this section.

- 1** Ethanol can be oxidised to ethanoic acid in a two-step process.



- 0 1** . **1** State a suitable oxidising agent and how you would ensure that the oxidation to ethanoic acid is complete.

**[4 marks]**

---

---

---

---

---

- 0 1** . **2** State the colour change that you would observe in this oxidation.

**[1 mark]**

---

- 0 1** . **3** The boiling points of the organic compounds in the reaction mixture are shown in **Table 1**.

**Table 1**

Compound	ethanol	ethanal	ethanoic acid
Boiling point / °C	78	21	118

Draw a diagram to illustrate the apparatus you could use to separate ethanal from a mixture of these three compounds.

**[4 marks]**

- 0 1** . **4** Use your knowledge of structure and bonding to explain why it is possible to separate ethanal in this way.

**[2 marks]**

---

---

---

---

---

**Question 1 continues on the next page**

- 0 1 . 5** Describe a chemical test for an aldehyde.  
In your answer, give the reagent, conditions and expected observation.

**[3 marks]**

---

---

---

---

---

---

- 0 1 . 6** Describe a chemical test for a carboxylic acid.  
In your answer, give the reagent and expected observation.

**[2 marks]**

---


---

---

---

---

- 2 Ethanol and ethanoic acid react reversibly to form ethyl ethanoate and water according to the equation:



A mixture of 4.80 g of ethanoic acid and 0.120 mol of ethanol is allowed to reach equilibrium at 20 °C.

- The equilibrium mixture is placed in a graduated flask and the volume made up to 250 cm<sup>3</sup> with distilled water.
- A 25.0 cm<sup>3</sup> sample of this equilibrium mixture is titrated with sodium hydroxide added from a burette.
- The ethanoic acid in this sample reacts with 4.00 cm<sup>3</sup> of 0.400 mol dm<sup>-3</sup> sodium hydroxide solution.

- |   |   |
|---|---|
| 0 | 2 |
|---|---|

 . 

1
---

 Calculate the value for  $K_c$  for the reaction of ethanoic acid and ethanol at 20 °C. Give your answer to the appropriate precision.

[7 marks]

$K_c =$  \_\_\_\_\_

**Question 2 continues on the next page**

A student obtained the titration results given in **Table 2**.

**Table 2**

	<b>Rough</b>	<b>1</b>	<b>2</b>	<b>3</b>
<b>Final burette reading / cm<sup>3</sup></b>	4.60	8.65	12.85	16.80
<b>Initial burette reading / cm<sup>3</sup></b>	0.10	4.65	8.65	12.85
<b>Titre / cm<sup>3</sup></b>				

**0 2** . **2** Complete **Table 2**.

[1 mark]

**0 2** . **3** Calculate the mean titre and justify your choice of titres.

[2 marks]

Calculation

Mean titre = \_\_\_\_\_ cm<sup>3</sup>

Justification \_\_\_\_\_

**0 2** . **4** Phenolphthalein is a suitable indicator for this titration.

State the colour change at the end point.

[1 mark]

**0 2** . **5** The error in the mean titre for this experiment is  $\pm 0.15 \text{ cm}^3$ .

Calculate the percentage error in this mean titre.

[1 mark]

Percentage error = \_\_\_\_\_ %

**0 2 . 6** Suggest how, using the same mass of ethanoic acid, the experiment could be improved to reduce the percentage error.

**[2 marks]**

---

---

---

---

---

**Turn over for the next question**

3

A peptide is hydrolysed to form a solution containing a mixture of amino acids. This mixture is then analysed by silica gel thin-layer chromatography (TLC) using a developing solvent. The individual amino acids are identified from their  $R_f$  values.

Part of the practical procedure is given below.

1. **Wearing plastic gloves to hold a TLC plate**, draw a pencil line 1.5 cm from the bottom of the plate.
2. Use a capillary tube to apply a very small drop of the solution of amino acids to the mid-point of the pencil line.
3. Allow the spot to dry completely.
4. In the developing tank, add the developing solvent to **a depth of not more than 1 cm**.
5. Place your TLC plate in the developing tank **and seal the tank with a lid**.
6. Allow the developing solvent to rise up the plate to at least  $\frac{3}{4}$  of its height.
7. Remove the plate and quickly mark the position of the solvent front with a pencil.
8. Allow the plate to dry **in a fume cupboard**.

**0 3** . **1** Parts of the procedure are in bold text.

Explain why these parts of the procedure are essential.

**[4 marks]**

---

---

---

---

---

---

---

---

---

---



- 0 3 . 2** Outline the steps needed to locate the positions of the amino acids on the TLC plate and to determine their  $R_f$  values.

**[4 marks]**

---

---

---

---

---

---

---

---

---

---

- 0 3 . 3** Explain why different amino acids have different  $R_f$  values.

**[2 marks]**

---

---

---


---

---



**0 4 . 2**

A buffer solution is made by adding 0.0600 mol of sodium hydroxide to a solution containing 0.100 mol of ethanedioic acid ( $\text{H}_2\text{C}_2\text{O}_4$ ). Assume that the sodium hydroxide reacts as shown in the following equation and that in this buffer solution, the ethanedioic acid behaves as a monoprotic acid.



Calculate a value for the pH of the buffer solution.  
Give your answer to 2 decimal places.

**[5 marks]**

pH = \_\_\_\_\_

**Question 4 continues on the next page**

**0 4 . 3** In a titration, the end point was reached when  $25.0 \text{ cm}^3$  of an acidified solution containing ethanedioic acid reacted with  $20.20 \text{ cm}^3$  of  $0.0200 \text{ mol dm}^{-3}$  potassium manganate(VII) solution.

Deduce an equation for the reaction that occurs and use it to calculate the original concentration of the ethanedioic acid solution.

**[4 marks]**

Equation \_\_\_\_\_

Calculation

Original concentration = \_\_\_\_\_  $\text{mol dm}^{-3}$

- 0 4 . 4** A sample of ethanedioic acid was treated with an excess of an unknown alcohol in the presence of a strong acid catalyst. The products of the reaction were separated and analysed in a time of flight (TOF) mass spectrometer. Two peaks were observed at  $m/z = 104$  and  $118$ .

Identify the compounds responsible for the two peaks.

**[2 marks]**

---

---

---

---

---

- 0 4 . 5** Outline how the TOF mass spectrometer is able to separate the two ions with different  $m/z$  values to give two peaks.

**[4 marks]**

---

---

---

---

---

---

---

---

**Turn over for the next question**

## Section B

Answer **all** questions in this section.

Only **one** answer per question is allowed.


For each answer completely fill in the circle alongside the appropriate answer.


CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown. 

If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown. 

**0 5**

Which change requires the largest amount of energy?

[1 mark]



**0 6**

A sample of 2.18 g of oxygen gas has a volume of 1870 cm<sup>3</sup> at a pressure of 101 kPa.

What is the temperature of the gas?

The gas constant is  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ .

[1 mark]

**A** 167 K ☐


**B** 334 K ☐

**C** 668 K ☐

**D** 334 000 K ☐

0	7
---	---

An ester is hydrolysed as shown by the following equation.



What is the percentage yield of RCOOH when 0.50 g of RCOOH ( $M_r = 100$ ) is obtained from 1.0 g of RCOOR' ( $M_r = 150$ )?

[1 mark]


- |          |     |                       |
|----------|-----|-----------------------|
| <b>A</b> | 33% | <input type="radio"/> |
| <b>B</b> | 50% | <input type="radio"/> |
| <b>C</b> | 67% | <input type="radio"/> |
| <b>D</b> | 75% | <input type="radio"/> |

Turn over for the next question

**0 8**

**Figure 1** shows the Maxwell–Boltzmann energy distribution curves for molecules of a sample of a gas at two different temperatures.

**Figure 1**



Which letter represents the most probable energy of the molecules at the lower temperature?


**[1 mark]**

- A** ☐
- B** ☐
- C** ☐
- D** ☐



0 9

The rate equation for the hydrogenation of ethene



is  $\text{Rate} = k[\text{C}_2\text{H}_4][\text{H}_2]$

At a fixed temperature, the reaction mixture is compressed to triple the original pressure.


What is the factor by which the rate of reaction changes?

[1 mark]

- A**     6     ☐
- B**     9     ☐
- C**     12     ☐
- D**     27     ☐

1 0

When one mole of ammonia is heated to a given temperature, 50% of the compound dissociates and the following equilibrium is established.




What is the total number of moles of gas present in this equilibrium mixture?

[1 mark]

- A**     1.5     ☐
- B**     2.0     ☐
- C**     2.5     ☐
- D**     3.0     ☐

1 1

Which change would alter the value of the equilibrium constant ( $K_c$ ) for this reaction?



[1 mark]

- A**    Increasing the concentration of sulfur trioxide.     ☐
- B**    Increasing the concentration of sulfur dioxide.     ☐
- C**    Increasing the temperature.     ☐
- D**    Adding a catalyst.     ☐

**1 2**

What is the pH of a  $0.020 \text{ mol dm}^{-3}$  solution of a diprotic acid which is completely dissociated?

**[1 mark]**

**A** 1.00 ☐

**B** 1.40 ☐

**C** 1.70 ☐

**D** 4.00 ☐

**1 3**

Equal volumes of equimolar solutions of the following pairs of substances are mixed.

Which pair makes a buffer solution?

**[1 mark]**

**A** Ammonia and ammonium chloride. ☐

**B** Hydrochloric acid and sodium hydroxide. ☐

**C** Hydrochloric acid and sodium chloride. ☐

**D** Ethanoic acid and methanoic acid. ☐

**1 4**

In which compound is the Group 7 element in its lowest oxidation state?

**[1 mark]**

**A** NaF ☐

**B** NaClO<sub>4</sub> ☐

**C** HOBr ☐

**D** KIO<sub>3</sub> ☐

**1 5**

In which reaction is hydrogen acting as an oxidising agent?

**[1 mark]**

**A**  $\text{Cl}_2 + \text{H}_2 \longrightarrow 2\text{HCl}$  ☐

**B**  $(\text{CH}_3)_2\text{CO} + \text{H}_2 \longrightarrow (\text{CH}_3)_2\text{CHOH}$  ☐




**C**  $\text{N}_2 + 3\text{H}_2 \longrightarrow 2\text{NH}_3$  ☐

**D**  $2\text{Na} + \text{H}_2 \longrightarrow 2\text{NaH}$  ☐

1 6


In which reaction is the metal oxidised?

[1 mark]



1 7

The following cell has an EMF of +0.46 V.



Which statement is correct about the operation of the cell?

[1 mark]

**A** Metallic copper is oxidised by  $\text{Ag}^{+}$  ions. ☐


**B** The silver electrode has a negative polarity. ☐

**C** The silver electrode gradually dissolves to form  $\text{Ag}^{+}$  ions. ☐

**D** Electrons flow from the silver electrode to the copper electrode via an external circuit. ☐

1 8

In an experiment to identify a Group 2 metal (X), 0.102 g of X reacts with an excess of aqueous hydrochloric acid according to the following equation.



The volume of hydrogen gas given off is  $65 \text{ cm}^3$  at 99 kPa pressure and 303 K. The gas constant is  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ .

Which is X?

[1 mark]

**A** Barium ☐

**B** Calcium ☐

**C** Magnesium ☐

**D** Strontium ☐

**1 9**

What forms when a solution of sodium carbonate is added to a solution of gallium(III) nitrate?

**[1 mark]**

**A** A white precipitate of gallium(III) carbonate.

☐

**B** A white precipitate of gallium(III) hydroxide.

☐

**C** A white precipitate of gallium(III) carbonate and bubbles of carbon dioxide.

☐

**D** A white precipitate of gallium(III) hydroxide and bubbles of carbon dioxide.

☐**2 0**

Which compound gives a colourless solution when an excess of dilute aqueous ammonia is added?

**[1 mark]**

**A**  $\text{MgCl}_2$

☐

**B**  $\text{AgCl}$

☐

**C**  $\text{CuCl}_2$

☐

**D**  $\text{AlCl}_3$

☐**2 1**

What is the final species produced when an excess of aqueous ammonia is added to aqueous aluminium chloride?

**[1 mark]**

**A**  $[\text{Al}(\text{NH}_3)_6]^{3+}$

☐

**B**  $[\text{Al}(\text{OH})_3(\text{H}_2\text{O})_3]$

☐

**C**  $[\text{Al}(\text{OH})_4(\text{H}_2\text{O})_2]^-$


☐

**D**  $[\text{Al}(\text{OH})(\text{H}_2\text{O})_5]^{2+}$

☐

2 2

The following equation represents the oxidation of vanadium(IV) ions by manganate(VII) ions in acid solution.



What volume of  $0.020 \text{ mol dm}^{-3}$   $\text{KMnO}_4$  solution is required to oxidise completely a solution containing  $0.010 \text{ mol}$  of vanadium(IV) ions?

[1 mark]

- A     $10 \text{ cm}^3$     ☐
- B     $25 \text{ cm}^3$     ☐
- C     $50 \text{ cm}^3$     ☐
- D     $100 \text{ cm}^3$     ☐

2 3

How many isomers have the molecular formula  $\text{C}_5\text{H}_{12}$ ?

[1 mark]

- A    2    ☐
- B    3    ☐
- C    4    ☐
- D    5    ☐

2 4

Which molecule is **not** produced when ethane reacts with bromine in the presence of ultraviolet light?

[1 mark]

- A     $\text{C}_2\text{H}_4\text{Br}_2$     ☐
- B     $\text{HBr}$     ☐
- C     $\text{H}_2$     ☐
- D     $\text{C}_4\text{H}_{10}$     ☐




**2 5**How many structural isomers have the molecular formula  $C_4H_9Br$ ?**[1 mark]****A** 2 ☐**B** 3 ☐**C** 4 ☐**D** 5 ☐**2 6**What is the major product of the reaction between but-1-ene and  $DBr$ ?  
(D is deuterium and represents  $^2H$ )**[1 mark]****A**  $CH_2DCH_2CH_2CH_2Br$  ☐**B**  $CH_2DCH_2CHBrCH_3$  ☐**C**  $CH_3CH_2CHBrCH_2D$  ☐**D**  $CH_3CH_2CHDCH_2Br$  ☐**2 7**

Why are fluoroalkanes unreactive?




**[1 mark]****A** Fluorine is highly electronegative. ☐**B** The  $F^-$  ion is very stable. ☐**C** They are polar molecules. ☐**D** The  $C-F$  bond is very strong. ☐**2 8**Which alcohol could **not** be produced by the reduction of an aldehyde or a ketone?**[1 mark]****A** 2-methylbutan-1-ol ☐**B** 2-methylbutan-2-ol ☐**C** 3-methylbutan-1-ol ☐**D** 3-methylbutan-2-ol ☐

**2 9**



Which compound forms optically active compounds on reduction?

**[1 mark]****3 0**How many secondary amines have the molecular formula  $\text{C}_4\text{H}_{11}\text{N}$ ?**[1 mark]****3 1**

Which compound has the highest boiling point?

**[1 mark]****3 2**

Which compound can polymerise by reaction with itself?

**[1 mark]**


3 3

A drug is designed to simulate one of the following molecules that absorbs onto the active site of an enzyme.


Which molecule requires the design of an optically active drug?

[1 mark]


A


☐


B


☐

C


☐

D


☐

3 4

Which amine has only **three** peaks in its proton NMR spectrum?

[1 mark]

A Methylamine

☐

B Trimethylamine

☐

C Diethylamine

☐

D Propylamine

☐

**END OF QUESTIONS**