



Cambridge International Examinations
Cambridge Pre-U Certificate

FURTHER MATHEMATICS (PRINCIPAL)

9795/01

Paper 1 Further Pure Mathematics

For Examination from 2016

SPECIMEN MARK SCHEME

3 hours

MAXIMUM MARK: 120

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 3 Pre-U Certificate.

This document consists of **10** printed pages.

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.

The following abbreviations may be used in a mark scheme:

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- CAO Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
- aef Any equivalent form
- art Answers rounding to
- cwo Correct working only (emphasising that there must be no incorrect working in the solution)
- ft Follow through from previous error is allowed
- o.e. Or equivalent


1		$\sum_{r=1}^n (r^2 - r + 1) = \sum_{r=1}^n r^2 - \sum_{r=1}^n r + \sum_{r=1}^n 1$ <p>Splitting summation and use of given results</p> $= \frac{1}{6}n(n+1)(2n+1) - \frac{1}{2}n(n+1) + n$ <p>1st for Σr^2; 2nd for Σr & $\Sigma 1 = n$</p> $= \frac{1}{3}n(n^2 + 2)$ <p>legitimately</p>	<p>M1</p> <p>B1 B1</p> <p>A1</p>
2		$A = k \int (\sin \theta + \cos \theta)^2 d\theta$ <p>including squaring attempt; ignore limits and $k \neq \frac{1}{2}$</p> $= \frac{1}{2} \int (1 + \sin 2\theta) d\theta$ <p>for use of the double-angle formula</p> <p>OR integration of $\sin \theta \cos \theta$ as $k \sin^2 \theta$ or $k \cos^2 \theta$</p> $= \frac{1}{2} \left[\theta - \frac{1}{2} \cos 2\theta \right]_0^{\pi/2}$ <p>ft (constants only) in the integration;</p> <p>MUST be 2 separate terms</p> $= \frac{1}{4} \pi + \frac{1}{2}$	<p>M1</p> <p>B1</p> <p>A1</p> <p>A1</p>
3	<p>(i)</p> <p>(ii)</p>	$y = (\sinh x)^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \frac{1}{2} (\sinh x)^{-\frac{1}{2}} \cdot \cosh x$ <p>OR $y^2 = \sinh x \Rightarrow 2y \frac{dy}{dx} = \cosh x$</p> $= \frac{\sqrt{1+y^4}}{2y}$ $\int \frac{2y}{\sqrt{1+y^4}} dy = \int 1 dx$ <p>By separating variables in (i)'s answer</p> $\Rightarrow x = \int \frac{2y}{\sqrt{1+y^4}} dy$ <p>But $x = \sinh^{-1} y^2$ so $\int \frac{2t}{\sqrt{1+t^4}} dx = \sinh^{-1}(t^2) + C$ condone missing “+ C”</p> <p>ALT.1 Set $t^2 = \sinh \theta$, $2t dt = \cosh \theta d\theta$ M1 Full substitution</p> $\int \frac{2t}{\sqrt{1+t^4}} dt = \int \frac{\cosh \theta}{\sqrt{1 + \sinh^2 \theta}} d\theta$ <p>A1 $= \int 1 \cdot d\theta = \theta = \sinh^{-1}(t^2)$ A1</p> <p>ALT.2 Set $t^2 = \tan \theta$, $2t dt = \sec^2 \theta d\theta$ M1 Full substitution</p> $\int \frac{2t}{\sqrt{1+t^4}} dt = \int \frac{\sec^2 \theta}{\sqrt{1 + \tan^2 \theta}} d\theta$ <p>A1 $= \int \sec \theta \cdot d\theta$</p> $= \ln \sec \theta + \tan \theta = \ln t^2 + \sqrt{1+t^4} $ <p>A1</p>	<p>M1 A1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p>


4	(i)	$y = \frac{x+1}{x^2+3} \Rightarrow y.x^2 - x + (3y-1) = 0$ Creating a quadratic in x For real x , $1 - 4y(3y-1) \geq 0$ Considering the discriminant $12y^2 - 4y - 1 \leq 0$ Creating a quadratic inequality For real x , $(6y+1)(2y-1) \leq 0$ Factorising/solving a 3-term quadratic $-\frac{1}{6} \leq y \leq \frac{1}{2}$ CAO (ii) $y = \frac{1}{2}$ substituted back $\Rightarrow \frac{1}{2}(x^2 - 2x + 1) = 0 \Rightarrow x = 1$ [$y = \frac{1}{2}$] $y = -\frac{1}{6}$ substituted back $\Rightarrow -\frac{1}{6}(x^2 + 6x + 9) = 0 \Rightarrow x = -3$ [$y = -\frac{1}{6}$]	M1 M1 M1 M1 A1 M1 A1 M1 A1
5	(i) (a) (b) (ii)	$\begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$ $\begin{pmatrix} \cos \beta & \sin \beta \\ \sin \beta & -\cos \beta \end{pmatrix}$ $\begin{pmatrix} \cos \phi & \sin \phi \\ \sin \phi & -\cos \phi \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$ multiplication of 2 reflection matrices. Correct order. $= \begin{pmatrix} \cos \phi \cos \theta + \sin \phi \sin \theta & \cos \phi \sin \theta - \sin \phi \cos \theta \\ \sin \phi \cos \theta - \cos \phi \sin \theta & \cos \phi \cos \theta + \sin \phi \sin \theta \end{pmatrix}$ $= \begin{pmatrix} \cos(\phi - \theta) & -\sin(\phi - \theta) \\ \sin(\phi - \theta) & \cos(\phi - \theta) \end{pmatrix}$ Use of the addition formulae; correctly done ... giving a Rotation (about O) through $(\phi - \theta)$	B1 B1 M1 M1 M1 A1 M1 A1
6	(i) (ii) (iii)	Possible orders are 1, 2, 3, 4, 6 & 12 By <i>Lagrange's Theorem</i> , the order of an element divides the order of the group (since the order of an element \equiv the order of the subgroup generated by that element) E.g. $y = xyx \Rightarrow y \cdot x^2y = xyx \cdot x^2y$ by ③ $= xy \cdot x^3 \cdot y = xy \cdot y^2 \cdot y$ by ② $= x \cdot y^4 = x \cdot (y^2)^2$ [by ②] $= x \cdot (x^3)^2 = x \cdot e$ by ① 2 M's for first, correct uses of 2 different conditions; the A for the 3 rd condition used to clinch the result. Proving G not abelian: [e.g. by $xyx = y$ but $x^2 \neq e$] $\Rightarrow G$ not cyclic OR establishing a contradiction	B1 B1 M1 M1 A1 B1 B1

7	(i)	$\cos 4\theta + i \sin 4\theta = (c + is)^4$ Use of <i>de Moivre's Theorem</i> $= c^4 + 4c^3 \cdot is + 6c^2 \cdot i^2 s^2 + 4c \cdot i^3 s^3 + i^4 s^4$ Binomial expansion attempted $\cos 4\theta = c^4 - 6c^2 s^2 + s^4$ and $\sin 4\theta = 4c^3 s - 4cs^3$ Equating Re & Im parts $\tan 4\theta = \frac{\sin 4\theta}{\cos 4\theta} = \frac{4c^3 s - 4cs^3}{c^4 - 6c^2 s^2 + s^4}$ Dividing throughout by c^4 to get $\frac{4t - 4t^3}{1 - 6t^2 + t^4}$ legitimately	M1 M1 M1 M1 A1
	(ii)	$t = \frac{1}{5} \Rightarrow \tan 4\theta = \frac{120}{119}$ $\tan\left(\frac{1}{4}\pi + \tan^{-1} \frac{1}{239}\right) = \frac{1 + \frac{1}{239}}{1 - \frac{1}{239}} = \frac{120}{119}$ Noting that this is $\tan(4\tan^{-1} \frac{1}{5})$ so that $4\tan^{-1} \frac{1}{5} = \frac{1}{4}\pi + \tan^{-1} \frac{1}{239}$	B1 M1 A1 A1
	(i)	Substituting $x = 1$, $f(1) = 2$ and $f'(1) = 3$ into (*) $\Rightarrow f''(1) = 5$	M1 A1
	(ii)	$\{x^2 f'''(x) + 2xf''(x)\} + \{(2x-1)f''(x) + 2f'(x)\} - 2f'(x) = 3e^{x-1}$ Product Rule used twice; at least one bracket correct Substituting $x = 1$, $f'(1) = 3$ and $f''(1) = 5$ into this $\Rightarrow f'''(1) = -12$ ft their $f''(1)$	M1 A1 M1 A1
	(iii)	$f(x) = f(1) + f'(1)(x-1) + \frac{1}{2}f''(1)(x-1)^2 + \frac{1}{6}f'''(1)(x-1)^3 + \dots$ Use of the Taylor series $= 2 + 3(x-1) + \frac{5}{2}(x-1)^2 - 2(x-1)^3 + \dots$ 1 st two terms CAO; 2 nd two terms ft (i) & (ii)'s answers	M1 A1 A1
	(iv)	Substituting $x = 1.1 \Rightarrow f(1.1) \approx 2.323$ to 3d.p. CAO	M1 A1
9	(i)	$\frac{dy}{dx} + y = 3xy^4$ is a <i>Bernoulli (differential) equation</i> $u = \frac{1}{y^3} \Rightarrow \frac{du}{dx} = -\frac{3}{y^4} \times \frac{dy}{dx}$ Then $\frac{dy}{dx} + y = 3xy^4$ becomes $-\frac{3}{y^4} \times \frac{dy}{dx} - \frac{3}{y^3} = -9x \Rightarrow \frac{du}{dx} - 3u = -9x$ AG	B1 M1 A1

	(ii)	<p>Method 1</p> <p>IF is e^{-3x}</p> $\Rightarrow ue^{-3x} = \int -9xe^{-3x} dx$ $= 3xe^{-3x} - \int 3e^{-3x} dx$ <p>Use of “parts”</p> $= (3x + 1)e^{-3x} + C$ <p>General solution is $u = 3x + 1 + Ce^{3x}$ ft</p> $\Rightarrow y^3 = \frac{1}{3x + 1 + Ce^{3x}}$ ft <p>Using $x = 0, y = \frac{1}{2}$ to find C $C = 7$ or $y^3 = \frac{1}{3x + 1 + 7e^{3x}}$</p> <p>Method 2</p> <p>Auxiliary equation $m - 3 = 0 \Rightarrow u_C = Ae^{3x}$ is the complementary function</p> <p>For particular integral try $u_P = ax + b$, $u_P' = a$</p> <p>Substituting $u_P = ax + b$ and $u_P' = a$ into the d.e. and comparing terms</p> $a - 3ax - 3b = -9x \Rightarrow a = 3, b = 1 \quad \text{i.e. } u_P = 3x + 1$ <p>General solution is $u = 3x + 1 + Ae^{3x}$ ft particular integral + complementary function provided particular integral has no arbitrary constants and complementary function has one</p> $\Rightarrow y^3 = \frac{1}{3x + 1 + Ae^{3x}}$ ft <p>Using $x = 0, y = \frac{1}{2}$ to find A $A = 7$ or $y^3 = \frac{1}{3x + 1 + 7e^{3x}}$</p>	<p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1 A1</p>
10	(i)	<p>Substituting $\begin{pmatrix} 1 + 3\lambda \\ -3 + 4\lambda \\ 2 + 6\lambda \end{pmatrix}$ into plane equation; i.e. $\begin{pmatrix} 1 + 3\lambda \\ -3 + 4\lambda \\ 2 + 6\lambda \end{pmatrix} \bullet \begin{pmatrix} 2 \\ -6 \\ 3 \end{pmatrix} = k$</p> <p>OR any point on line (since “given”)</p> $k = 2 + 6\lambda + 18 - 24\lambda + 6 + 18\lambda = 26$	<p>M1</p> <p>A1</p>

	<p>(ii) Working with vector $\begin{pmatrix} 10+2m \\ 2-6m \\ 3m-43 \end{pmatrix}$.</p> <p>Substituting into the plane equation: $\begin{pmatrix} 10+2m \\ 2-6m \\ 3m-43 \end{pmatrix} \bullet \begin{pmatrix} 2 \\ -6 \\ 3 \end{pmatrix} = k$</p> <p>Solving a linear equation in m: $20 + 4m - 12 + 36m + 9m - 129 = 26$</p> <p>$m = 3 \Rightarrow Q = (16, -16, -34)$</p> <p>Shortest distance is $m \left \begin{pmatrix} 2 \\ -6 \\ 3 \end{pmatrix} \right = 21$ or $PQ = \sqrt{6^2 + 18^2 + 9^2} = 21$</p> <p>(iii) Finding 3 points in the plane: e.g. $A(1, -3, 2)$, $B(4, 1, 8)$, $C(10, 2, -43)$</p> <p>Then 2 vectors in (\parallel to) plane: e.g. $\overrightarrow{AB} = \begin{pmatrix} 3 \\ 4 \\ 6 \end{pmatrix}$, $\overrightarrow{AC} = \begin{pmatrix} 9 \\ 5 \\ -45 \end{pmatrix}$, $\overrightarrow{BC} = \begin{pmatrix} 6 \\ 1 \\ -51 \end{pmatrix}$</p> <p>OR B1 B1 for any two vectors in the plane</p> <p>Vector product of any two of these to get normal to plane: $\begin{pmatrix} 10 \\ -9 \\ 1 \end{pmatrix}$</p> <p>(any non-zero multiple)</p> <p>$d = \begin{pmatrix} 10 \\ -9 \\ 1 \end{pmatrix} \bullet (\text{any position vector}) = \begin{pmatrix} 10 \\ -9 \\ 1 \end{pmatrix} \bullet \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix}$ e.g. $= 39$</p> <p>$\Rightarrow 10x - 9y + z = 39$ CAO (aef)</p> <p>ALTERNATE SOLUTION</p> <p>$ax + by + cz = d$ contains $\begin{pmatrix} 1+3\lambda \\ -3+4\lambda \\ 2+6\lambda \end{pmatrix}$ and $\begin{pmatrix} 10 \\ 2 \\ -43 \end{pmatrix}$</p> <p>... so $a + 3a\lambda + 4b\lambda - 3b + 2c + 6c\lambda = d$ and $10a + 2b - 43c = d$</p> <p>Then $a - 3b + 2c = d$ and $3a + 4b + 6c = 0$ (λ terms) i.e. equating terms</p> <p>Eliminating (e.g.) c from 1st two equations $\Rightarrow 9a + 10b = 0$</p> <p>Choosing $a = 10, b = -9 \Rightarrow c = 1$ and $d = 39$ i.e. $10x - 9y + z = 39$ CAO</p>	<p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1 B1</p> <p>M1</p> <p>M1</p> <p>M1 A1</p>
--	---	---

11	<p>(i) $w = \sqrt{(\sqrt{3}-1)^2 + (\sqrt{3}+1)^2} = \sqrt{4-2\sqrt{3}+4+2\sqrt{3}} = \sqrt{8} \text{ or } 2\sqrt{2}$</p> <p>$\arg(w) = \tan^{-1}\left(\frac{\sqrt{3}+1}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}\right) = \tan^{-1}(2+\sqrt{3}) = \frac{5}{12}\pi$</p> <p>(ii) (a) $z^3 = \left(2\sqrt{2}, \frac{5}{12}\pi\right), \left(2\sqrt{2}, \frac{29}{12}\pi\right), \left(2\sqrt{2}, -\frac{19}{12}\pi \text{ or } \frac{53}{12}\pi\right)$</p> <p>$\sqrt[3]{ w }; \frac{\arg(w)}{3}$ These method marks can be earned for just the first root</p> <p>$\Rightarrow z = \left(\sqrt{2}, \frac{5}{36}\pi\right), \left(\sqrt{2}, \frac{29}{36}\pi\right), \left(\sqrt{2}, -\frac{19}{36}\pi\right)$ A marks for the 2nd & 3rd roots: $r e^{i\theta}$ forms equally acceptable</p> <p>(b) z_1, z_2, z_3 the roots of $z^3 - 0z^2 + 0z - w = 0$ $\Rightarrow z_1 z_2 z_3 = w = (\sqrt{3}-1) + i(\sqrt{3}+1)$ ALT. Multiplying the 3 roots together in any form</p> <p>(c)</p> <div style="display: flex; align-items: center;">  <div style="margin-left: 20px;"> <p>Three points in approx. correct places</p> <p>All equally spaced around a circle, centre O, radius $\sqrt{2}$ (Explained that Δ_1 equilateral)</p> <p>$l = 2 \times \sqrt{2} \sin\left(\frac{1}{2} \times \frac{2}{3}\pi\right) = \sqrt{6}$ or by the <i>Cosine Rule</i></p> </div> </div> <p>(d) $k = \exp\left\{-i, \frac{5}{36}\pi\right\} \text{ or } \exp\left\{-i, \frac{29}{36}\pi\right\} \text{ or } \exp\left\{i, \frac{19}{36}\pi\right\}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1 M1</p> <p>A1 A1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>B1</p>
----	--	---

12	(i)	$I_n = \int_0^3 x^{n-1} (x\sqrt{16+x^2}) \, dx$ <p>Correct splitting <i>and</i> use of parts</p> $= \left[x^{n-1} \cdot \frac{(16+x^2)^{3/2}}{3} \right]_0^3 - \int_0^3 (n-1)x^{n-2} \frac{(16+x^2)^{3/2}}{3} \, dx$ $= 3^{n-2} \cdot 125 - \left(\frac{n-1}{3} \right) \int_0^3 x^{n-2} (16+x^2) \sqrt{16+x^2} \, dx$ <p>Method to get 2nd integral of correct form</p> $= 3^{n-2} \cdot 125 - \left(\frac{n-1}{3} \right) \{16I_{n-2} + I_n\} \quad [\text{i.e. reverting to } I\text{'s in 2nd integral ft}]$ $\Rightarrow 3I_n = 3^{n-1} \cdot 125 - 16(n-1)I_{n-2} - (n-1)I_n$ <p>Collecting up I_ns</p> $(n+2)I_n = 125 \times 3^{n-1} - 16(n-1)I_{n-2} \quad \mathbf{AG}$	M1
			A1
			M1
			M1
			A1
	(ii) (a)	 <p>Spiral (with r increasing)</p> <p>From O to just short of $\theta = \pi$</p>	B1
			B1
	(b)	$r = \frac{1}{4}\theta^4 \Rightarrow \frac{dr}{d\theta} = \theta^3 \quad \text{and} \quad r^2 + \left(\frac{dr}{d\theta} \right)^2 = \frac{1}{16}\theta^8 + \theta^6$ $L = \int_0^3 \frac{1}{4}\theta^3 \sqrt{16+\theta^2} \, d\theta \quad (= \frac{1}{4}I_3)$ <p>Now $I_1 = \left[\frac{1}{3}(16+x^2)^{3/2} \right]_0^3 = \frac{61}{3}$</p> <p>and $5I_3 = 125 \times 9 - 16 \times 2 \left(\frac{61}{3} \right) = \frac{1423}{3}$ or $474\frac{1}{3}$ Use of given reduction formula</p> <p>so that $L = \frac{1}{20} \times \frac{1423}{3} = \frac{1423}{60}$ or $23\frac{43}{60}$ or awrt 23.7 ft only from suitable $k I_3$</p>	M1 A1
			M1 A1
			B1
			M1
			A1

13	<p>Base-line case: for $n = 5$, $13579 R_5 = 1508\ 7\ 6269$ contains a string of $(5 - 4 = 1)$ 7s</p> <p>$13579 R_6 = 1508776269$, $13579 R_7 = 15087776269$, etc. or form of 1st & last 4 digits</p> <p>Assume that, for some $k \geq 5$, $13579 R_k = 1508 \frac{77\dots7}{(k-4)\ 7\text{'s}} 6269$. Induction hypothesis</p> <p>Then, for $n = k + 1$,</p> $13579 R_{k+1} = 13579(10R_k + 1)$ <p>Give the M mark for the key observation that $R_{k+1} = 10R_k + 1$ or $10^k + R_k$, even if not subsequently used.</p> $ \begin{array}{r} = 1508 \frac{77\dots7}{(k-4)\ 7\text{'s}} 62690 \\ + 13579 \\ = 1508 \frac{77\dots7}{(k-4+1)\ 7\text{'s}} 76269 \end{array} $ <p>which contains a string of $(k - 4 + 1)$ 7s, as required. Proof follows by induction (usual round-up).</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>
----	--	---